

# Virtual Patient Simulation is Highly Effective at Improving Guideline-Based Management of Hyperlipidemia

Presented at ACC 2020 - 69th Annual Scientific Session & Expo of the American College of Cardiology 28th - 30th March, 2020 Chicago, IL, USA


AMY LARKIN, PHARMD; DAVID ANDERSON; MARTIN WARTERS, CHSE; ALLIE ARMAGAN, PHARMD, Medscape Education, New York, NY

## PURPOSE

We sought to determine if an online, virtual patient simulation (VPS)based continuing medical education (CME) intervention could improve performance of cardiologists and primary care physicians (PCPs) related to guideline-based management of patients with hyperlipidemia.

### METHODS

- The intervention comprised 2 patients presenting in a VPS platform that allows learners to order lab tests, make diagnoses, and prescribe treatments supported by an extensive database of diagnostic and treatment possibilities matching the scope and depth of actual practice
- Learner clinical decisions, entered using open field entries, were analyzed using a sophisticated decision engine
  - Tailored clinical guidance (CG), based on current evidence and expert recommendation, was provided following each decision
  - Learners were given the opportunity to modify their decisions after receiving CG
- Each user's decisions were collected post-CG and compared with that user's pre-CG data using a 2-tailed, paired t-test
- Data are reflective of learners who participated in the VPS from April 29, 2019, through July 25, 2019




## RESULTS


#### Patient Simulation Cases


BMI: 31

Overall, significant improvements were demonstrated by both cardiologists and PCPs related to comprehensive management of patients with hyperlipidemia.

In total, 105 cardiologists and 375 PCPs were included. Significant improvements (reported as absolute differences) were observed after CG.







## CONCLUSIONS

- VPS that immerses and engages specialists in an authentic and practical learning experience can improve evidence-based clinical decisions related to guideline-based management of hyperlipidemia
- Persistent educational gaps were uncovered related to:
- Order TIMI risk score for secondary prevention
- Order Lipoprotein (a) test
- Initiate a PCSK9 inhibitors or other nonstatin therapy when appropriate

#### DISCLOSURE

The author has nothing to disclose.

#### ACKNOWLEDGEMENTS

This CME activity was supported by an independent educational grant from Amgen.

For more information, contact Amy Larkin, PharmD, Senior Director, Clinical Strategy, Medscape, LLC, at alarkin@medscape.net.



Scan here to view this poster online.